Brain Research, 531 (1990) 358–359
Elsevier

Author Index

Aebischer, P., Guénard, V. and Valentini, R.F., The morphology of regenerating peripheral nerves is modulated by the surface microgeometry of polymeric guidance channels, 211
April, M., see Heyes, M.P., 148
Aulerich, R.J., see Tanaka Jr., D., 294

Barry, D.I., see Hansen, A., 307
Bartolome, J.V., see Lorber, B.A., 329
Bass, A., see Fluet, A., 312
Bates, J.N., see Meller, S.T., 342
Ben-Jonathan, N., see Burgett, R.A., 143
Bienvenu, O.J., see Goeders, N.E., 322
Blackmore, D., see Heyes, M.P., 148
Bloom, F.E., see Giovannelli, L., 299
Boer, H.H., see Leung, M.K., 66
Bolwig, T.G., see Hansen, A., 307
Boyce, S., see Robertson, R.G., 95
Brody, M.J., see Meller, S.T., 342
Buchanan, J., see Simpson III, R.E., 136
Burgett, R.A., Garris, P.A. and Ben-Jonathan, N., Estradiol-induced prolactinomas: differential effects on dopamine in posterior pituitary and median eminence, 143
Bursian, S.J., see Tanaka Jr., D., 294

Carenzi, A., see Pezzoli, G., 355
Carlen, P.L., see Mody, I., 88
Clarke, C.A., see Robertson, R.G., 95
Clarke, R.W., see Taylor, J.S., 263
Cox, B.M., see Izenwasser, S., 338
Crain, S.M., see Shen, K.-F., 1
Crossman, A.R., see Roberston, R.G., 95

Dalsass, M. and Siegel, A., Opioid peptide regulation of neurons in the bed nucleus of the stria terminalis: a microiontophoretic study, 346
Dam, T.-V., Martinelli, B. and Quirion, R., Autoradiographic distribution of brain neurokinin-1/substance P receptors using a highly selective ligand [^{3}H]-[Sar-^{9},Met-(O_2)^{11}]-substance P, 333
Davies, P., see Vincent, I.J., 127
Dawson Jr., R., see Ulshafer, R.J., 350
De Graaf, F., see Van Asselt, E., 25
DeLanney, L.E., see Irwin, I., 242
De Souza, E.B., see Goeders, N.E., 322
Diegenbach, P.C., see Van Asselt, E., 25
Di Monte, D.A., see Irwin, I., 242
Dragunow, M., see Gunn, A.J., 105
Drucker-Colín, R., see Pacheco-Cano, M.T., 290
Dučić, I., see Polc, P., 286
Ducis, I., Norenberg, L.-O.B. and Norenberg, M.D., The benzodiazepine receptor in cultured astrocytes from genetically epilepsy-prone rats, 318
Dyck, P.J., see Yasaki, S., 8

Eckhaus, M., see Heyes, M.P., 148
Eneroth, P., see Von Euler, G., 253

Faull, R.L.M., see Gunn, A.J., 105
Finnegan, K.T., see Irwin, I., 242
Fluet, A. and Bass, A., Sexual dimorphisms in the vocal control system of a teleost fish: ultrastructure of neuromuscular junctions, 312
Ford, T.W., see Taylor, J.S., 263
Forno, L.S., see Irwin, I., 242
Franceschini, N., see Picaud, S., 117
Freitag, S.K., see Lorber, B.A., 329
Fuxe, K., see Von Euler, G., 253

García-Hernández, F., see Pacheco-Cano, M.T., 290
Garris, P.A., see Burgett, R.A., 143
Gebhart, G.F., see Meller, S.T., 342
Giovanelli, L., Shiromani, P.J., Jirikowski, G.F. and Bloom, F.E., Oxytocin neurons in the rat hypothalamus exhibit c-*fos* immunoreactivity upon osmotic stress, 299
Gluckman, P.D., see Gunn, A.J., 105
Goeders, N.E., Bienvenu, O.J. and De Souza, E.B., Chronic cocaine administration alters corticotropin-releasing factor receptors in the rat brain, 322
Gould, E., see Woolley, C.S., 225
Gravell, M., see Heyes, M.P., 148
Gruber, M.B., see Lebrun, C.J., 167
Guénard, V., see Aebischer, P., 211
Gunn, A.J., Dragunow, M., Faull, R.L.M. and Gluckman, P.D., Effects of hypoxia-ischemia and seizures on neuronal and glial-like c-*fos* protein levels in the infant rat, 105

Hansen, A., Jørgensen, O.S., Bolwig, T.G. and Barry, D.I., Hippocampal kindling alters the concentration of glial fibrillary acidic protein and other marker proteins in rat brain, 307
Hansen, L.A., see Masliah, E., 36
Harris, J., see Taylor, J.S., 263
Heuts, B., see Van Asselt, E., 25
Heyes, M.P., Gravell, M., London, W.T., Eckhaus, M., Vickers, J.H., Yergey, J.A., April, M., Blackmore, D. and Markey, S.P., Sustained increases in cerebrospinal fluid quinolinic acid concentrations in rhesus macaques (*Macaca mulatta*) naturally infected with simian retrovirus type-D, 148
Hiriart, M., see Pacheco-Cano, M.T., 290
Hökfelt, T., see Von Euler, G., 253
Horne, A.L., see Priestley, T., 183
Humlová, M. and Illnerová, H., Rate of reentrainment of circadian rhythms to advances of light-dark cycles may depend on ways of shifting the cycles, 304

Iimoto, D.S., see Masliah, E., 36
Illnerová, H., see Humlová, M., 304
Imai, S., see Onozuka, M., 276
Irwin, I., DeLanney, L.E., Forno, L.S., Finnegan, K.T., Di Monte, D.A. and Langston, J.W., The evolution of nigrostriatal neurochemical changes in the MPTP-treated squirrel monkey, 242
Isaac, L., Van Zandt O'Malley, T., Ristic, H. and Stewart, P., MK-801 blocks dynorphin A (1–13)-induced loss of the tail-flick reflex in the rat, 83
Ishikawa, T., see Yang, H., 280
Izenwasser, S. and Cox, B.M., Daily cocaine treatment produces a persistent reduction of [^{3}H]dopamine uptake in vitro in rat nucleus accumbens but not in striatum, 338

Jacobowitz, D.M., see Stanfield, B.B., 219
Jacobs, B.L., see Levine, E.S., 189
Jensen, M.S., see Nathan, T., 55
Jirikowski, G.F., see Giovanelli, L., 299
Jørgensen, O.S., see Hansen, A., 307

Kaneko, T., see Tashiro, Y., 159
Kemp, J.A., see Priestley, T., 183
Kikuchi, H., see Tashiro, Y., 159
Knight, R.T., see Singh, J., 45
Komisaruk, B.R., see Pacheco-Cano, M.T., 290
Kurumaji, A. and McCulloch, J., Effects of MK-801 upon local cerebral glucose utilisation in conscious rats following unilateral lesion of caudal entorhinal cortex, 72

Lambert, J.D.C., see Nathan, T., 55
Langston, J.W., see Irwin, I., 242
Lebrun, C.J., Gruber, M.G., Meister, M. and Unger, Th., Central vasopressin pretreatment sensitizes phosphoinositol hydrolysis in the rat septum, 167
Lehning, E.J., see Tanaka Jr., D., 294
Leung, M.K., Boer, H.H., Van Minnen, J., Lundy, J. and Stefano, G.B., Evidence for an enkephalinergic system in the nervous system of the pond snail, *Lymnaea stagnalis*, 66
Levine, E.S., Litto, W.J. and Jacobs, B.L., Activity of cat locus coeruleus noradrenergic neurons during the defense reaction, 189
Lewis, S.J., see Meller, S.T., 342
Litto, W.J., see Levine, E.S., 189
London, W.T., see Heyes, M.P., 148
Lorber, B.A., Freitag, S.K. and Bartolome, J.V., Effects of beta-endorphin on DNA synthesis in brain regions of preweanling rats, 329
Lowenstein, D.H., Simon, R.P. and Sharp, F.R., The pattern of 72-kDa heat shock protein-like immunoreactivity in the rat brain following fluorothyl-induced status epilepticus, 173
Lundy, J., see Leung, M.K., 66

Ma, R., see Wang, W., 196
MacDonald, J.F., see Mody, I., 88
Mariani, C.B., see Pezzoli, G., 355

Markey, S.P., see Heyes, M.P., 148
Martinelli, B., see Dam, T.-V., 333
Masliah, E., Iimoto, D.S., Saitoh, T., Hansen, L.A. and Terry, R.D., Increased immunoreactivity of brain spectrin in Alzheimer disease: a marker for synapse loss?, 36
Masotto, C., see Pezzoli, G., 355
McCulloch, J., see Kurumaji, A., 72
McEwen, B.S., see Woolley, C.S., 225
McKernan, R.M., see Priestley, T., 183
Meister, B., see Von Euler, G., 253
Meister, M., see Lebrun, C.J., 167
Meller, S.T., Lewis, S.J., Bates, J.N., Brody, M.J. and Gebhart, G.F., Is there a role for an endothelium-derived relaxing factor in nociception?, 342
Michels, K.M., Morin, L.P. and Moore, R.Y., $GABA_A$/benzodiazepine receptor localization in the circadian timing system, 16
Mizuno, N., see Tashiro, Y., 159
Mody, I., Reynolds, J.N., Salter, M.W., Carlen, P.L. and MacDonald, J.F., Kindling-induced epilepsy alters calcium currents in granule cells of rat hippocampal slices, 88
Moore, R.Y., see Michels, K.M., 16
Morin, L.P., see Michels, K.M., 16

Nagatsu, I., see Tashiro, Y., 159
Nathan, T., Jensen, M.S. and Lambert, J.D.C., $GABA_B$ receptors play a major role in paired-pulse facilitation in area CA1 of the rat hippocampus, 55
Norenberg, L.-O.B., see Ducis, I., 318
Norenberg, M.B., see Ducis, I., 318
Numan, R. and Quaranta Jr., J.R., Effects of medial septal lesions on operant delayed alternation in rats, 232

O'Neal, M., see Scremin, O.U., 203
Onozuka, M., Imai, S. and Ozono, S., The 70-kDa epileptic cortical protein elicits bursting activity accompanied by a reduction of outward current in *Euhadra* neurons which is inhibited by an antibody against this protein, 276
Ozono, S., see Onozuka, M., 267

Pacheco-Cano, M.T., García-Hernández, F., Hiriart, M., Komisaruk, B.R. and Drucker-Colín, R., Dibutyryl cAMP stimulates analgesia in rats bearing a ventricular adrenal medulla transplant, 290
Peichl, L., see Picaud, S., 117
Pettit, J.S., see Taylor, J.S., 263
Pezzoli, G., Ricciardi, S., Masotto, C., Mariani, C.B. and Carenzi, A., *n*-Hexane induces Parkinsonism in rodents, 355
Phillis, J.W., see Simpson III, R.E., 136
Picaud, S., Peichl, L. and Franceschini, N., Dye-induced 'photo-degeneration' and 'photo-permeabilization' of mammalian neurons in vivo, 117
Polc, P. and Dučić, I., Afferent stimulation frequency modulates GABAergic phenomena in the spinal cord: reversal by benzodiazepine antagonists, 286
Priestley, T., Horne, A.L., McKernan, R.M. and Kemp, J.A., The effect of NMDA receptor glycine site antagonists on hypoxia-induced neurodegeneration of rat cortical cell cultures, 183

Quaranta Jr., J.R., see Numan, R., 232
Quirion, R., see Dam, T.-V., 333

Reynolds, J.N., see Mody, I., 88
Ricciardi, S., see Pezzoli, G., 355
Richardson, J.S., see Tiong, A.H.K., 269
Ristic, H., see Isaac, L., 83
Robertson, R.G., Clarke, C.A., Boyce, S., Sambrook, M.A. and Crossman, A.R., The role of striatopallidal neurones utilizing gamma-aminobutyric acid in the pathophysiology of MPTP-induced parkinsonism in the primate: evidence from [^{3}H]flunitrazepam autoradiography, 95
Saitoh, T., see Masliah, E., 36
Salter, M.W., see Mody, I., 88
Sambrook, M.A., see Robertson, R.G., 95
Scremin, A.M.E., see Scremin, O.U., 203
Scremin, O.U., O'Neal, M. and Scremin, A.M.E., Glucose utilization and reflex activity of the transected rat spinal cord, 203
Sharp, F.R., see Lowenstein, D.H., 173
Shen, K.-F. and Crain, S.M., Cholera toxin-B subunit blocks excitatory effects of opioids on sensory neuron action potentials indicating that GM1 ganglioside may regulate G_s-linked opioid receptor functions, 1
Sherry, D.M., see Ulshafer, R.J., 350
Shiromani, P.J., see Giovannelli, L., 299
Siegel, A., see Dalsass, M., 346
Simon, R.P., see Lowenstein, D.H., 173
Simpson III, R.E., Phillis, J.W. and Buchannan, J., A comparison of cerebral blood flow during basal, hypotensive, hypoxic and hypercapnic conditions between normal and streptozotocin diabetic rats, 136
Singh, J. and Knight, R.T., Frontal lobe contribution to voluntary movements in humans, 45
Smit-Onel, M.J., see Van Asselt, E., 25
Stanfield, B.B. and Jacobowitz, D.M., Antibody to a soluble protein purified from brain selectively labels layers V corticofugal projection neurons in rat neocortex, 219
Stefano, G.B., see Leung, M.K., 66
Stewart, P., see Isaac, L., 83

Taché, Y., see Yang, H., 280
Tanaka, Jr., D., Bursian, S.J., Lehning, E.J. and Aulerich, R.J., Exposure to triphenyl phosphite results in widespread degeneration in the mammalian central nervous system, 294
Tashiro, Y., Kaneko, T., Nagatsu, I., Kikuchi, H. and Mizuno, N., Increase of tyrosine hydroxylase-like immunoreactive neurons in the nucleus accumbens and the olfactory bulb in the rat with the lesion in the ventral tegmental area of the midbrain, 159
Taylor, J.S., Pettit, J.S., Harris, J., Ford, T.W. and Clarke, R.W., Noxious stimulation of the toes evokes long-lasting, naloxone-reversible suppression of the sural-gastrocnemius reflex in the rabbit, 263
Terry, R.D., see Masliah, E., 36
Tiong, A.H.K. and Richardson, J.S., Differential effects of olfactory bulbectomy on β-adrenoceptors in rat amygdala, hippocampus and cerebral cortex, 269

Ulshafer, R.J., Sherry, D.M., Dawson Jr., R. and Wallace, D.R., Excitatory amino acid involvement in retinal degeneration, 350
Unger, Th., see Lebrun, C.J., 167
Valentini, R.F., see Aebischer, P., 211

Van Asselt, E., Van Raamsdonk, W., De Graaf, F., Smit-Onel, M.J., Diegenbach, P.C. and Heuts, B., Enzyme histochemical profiles of fish spinal motoneurons after cordotomy and axotomy of motor nerves, 25
Van Minnen, J., see Leung, M.K., 66
Van Raamsdonk, W., see Van Asselt, E., 25
Van Zandt, O'Malley, T., see Isaac, L., 83
Vickers, J.H., see Heyes, M.P., 148
Vincent, I.J. and Davies, P., Phosphorylation characteristics of the A68 protein in Alzheimer's disease, 127
Von Euler, G., Meister, B., Hökfelt, T., Eneroth, P. and Fuxe, K., Intraventricular injection of neurotensin reduces dopamine D_2 agonist binding in rat forebrain and intermediate lobe of the pituitary gland. Relationship to serum hormone levels and nerve terminal coexistence, 253

Wallace, D.R., see Ulshafer, R.J., 350
Wang, W. and Ma, R., The role of serotonin in non-cholinergic excitatory transmission in the guinea pig inferior mesenteric ganglion, 196
Woolley, C.S., Gould, E. and McEwen, B.S., Exposure to excess glucocorticoids alters dendritic morphology of adult hippocampal pyramidal neurons, 225

Yang, H., Ishikawa, T. and Taché, Y., Microinjection of TRH analogs into the raphe pallidus stimulates gastric acid secretion in the rat, 280
Yasaki, S. and Dyck, P.J., Duration and severity of hypoglycemia needed to induce neuropathy, 8
Yergey, J.A., see Heyes, M.P., 148